

CHARACTERIZATIONS AND ALGORITHMS FOR GENERALIZED COPS AND ROBBERS GAMES

ANTHONY BONATO AND GARY MACGILLIVRAY

ABSTRACT. We propose a definition of generalized Cops and Robbers games where there are two players, the Pursuer and the Evader, who may move via prescribed rules. If the Pursuer can ensure that the game enters into a fixed set of final positions, then the Pursuer wins; otherwise, the Evader wins. A relational characterization of the games where the Pursuer wins is provided. A precise formula is given for the length of the game, along with an algorithm for computing if the Pursuer has a winning strategy whose complexity is a function of the parameters of the game. For games where the position of one player does not affect the collection of moves available to the other, a vertex elimination ordering characterization is given for when the Pursuer has a winning strategy.

1. INTRODUCTION

Cops and Robbers, introduced over thirty years ago in [1, 16, 18, 19], is one example of a graph searching game that has received wide attention. The players—a set of *cops* and a *robber*—move alternately from vertex to vertex of a graph in a sequence of discrete time-steps. The cops attempt to capture the robber by occupying his vertex, while the robber is trying to avoid this situation forever. The consideration of Cops and Robbers gives rise to a number of deep results and problems in structural, algorithmic, and random graph theory; see the book [5] for a survey.

Many variations of Cops and Robbers are possible, for example, where different kinds of moves are allowed or forbidden, the cops or robber have more or less power (for example, the cops could win if they are sufficiently close to the robber). The game may also be played analogously on directed graphs or other relational structures. Our purpose here is not to survey all of these myriad variants, but rather, to offer a single framework for the analysis of *all* reasonably defined generalizations of Cops and Robbers which are perfect information games. We regard such games as being played between a Pursuer and an Evader, where the Pursuer’s objective is “capturing” the Evader, and the Evader’s objective is to avoid capture forever (hence, the Pursuer plays the role of the set of cops, and the Evader plays the role of the robber).

The paper is organized as follows. In Section 2, we present a general definition of generalized Cops and Robbers games and its associated bipartite state digraph, written \mathcal{D}_G . A labelling of the vertices of \mathcal{D}_G makes it possible to determine which player has a winning strategy, and also determines the length of the game. Section 3 provides a relational characterization of the games where the Pursuer has a winning strategy; see Theorem 1. This result generalizes the relational characterization of cop-win graphs from [17], and the one

2000 *Mathematics Subject Classification.* 05C57, 05C85.

Key words and phrases. Cops and Robbers, directed graphs, relational characterization, vertex elimination ordering.

The authors gratefully acknowledge support from NSERC.

given for k -cop-win graphs in [8, 15]. Our focus in the next section then shifts to those generalized Cops and Robbers games where the moves available to either side are not restricted by the position of the other side. For some of these games we are able to give a refinement of the relational characterization, as well as a characterization in terms of a vertex ordering of an associated product graph. We give an algorithm that determines if the Pursuer has a winning strategy. In the case of Cops and Robbers with k cops, this algorithm matches the time complexity of the known algorithms recognizing a winning strategy for the cops; see [3, 5, 8]. The characterizations and algorithms we provide have the advantage that they apply to contexts far more general than in the original Cops and Robbers, and to games played on different kinds of relational structures and with a variety of rules. In our final section, we consider an analogue of the cop number for generalized Cops and Robbers games.

2. GENERALIZED COPS AND ROBBERS GAMES

Intuition suggests that in a game generalized Cops and Robbers, each player should have his or her own position at any time in the game. The set of positions of the game should therefore consist of ordered pairs (p_P, q_E) , where p_P is the position of player the Pursuer, and q_E is the position of player the Evader. It may be that not every possible pair of positions is allowed or achievable. A move in a generalized Cops and Robbers game should involve a player possibly changing his own position, but not altering the position of his opponent. The *state* of the game consists of the current position of the game, and an identifier of which player is next to move; each state is an ordered pair $((p_P, q_E), X)$, where $X \in \{P, E\}$ indicates which player is next to move. Each player, on his turn, alters the state of the game by possibly changing his position, and changing the player who is next to move. We formalize these notions in the following definition, which has similarities to the definition of a combinatorial game.

A *generalized Cops and Robbers game* is any discrete-time process \mathcal{G} that satisfies the following rules:

- (1) There are two players named the *Pursuer* and the *Evader*.
- (2) There is perfect information.
- (3) There is a set \mathcal{P}_P of *allowed positions for the Pursuer* and a set \mathcal{P}_E of *allowed positions for the Evader*. The set of *positions of the game* is the subset $\mathcal{P} \subseteq \mathcal{P}_P \times \mathcal{P}_E$ of positions that can be achieved when moves are made according to the rules of the game. Similarly, the set of *states* of the game is the subset of $\mathcal{S} \subseteq \mathcal{P} \times \{P, E\}$ such that $((p_P, q_E), X) \in \mathcal{S}$ if the position (p_P, q_E) , with X next to move, can be achieved when moves are made according to the rules of the game.
- (4) For each state of the game and each player, there is a non-empty set of allowed moves. Each allowed move leaves the position of the other player unchanged. We use $\mathcal{A}_P(p_P, q_E)$ to denote the set of *allowed moves for the Pursuer* when the state of the game is $((p_P, q_E), L)$, and $\mathcal{A}_E(p_P, q_E)$ to denote the set of *allowed moves for the Evader* when the state of the game is $((p_P, q_E), R)$.
- (5) The rules of the game specify how it starts. That is, there is a set $\mathcal{I} \subseteq \mathcal{P}_P \times \mathcal{P}_E$ of *allowed start positions*. We define the set $\mathcal{I}_P = \{p_P : (p_P, q_E) \in \mathcal{I} \text{ for some } q_E \in \mathcal{P}_E\}$ and, for $p_P \in \mathcal{P}_P$, we define the set $\mathcal{I}_E(p_P) = \{q_E \in \mathcal{P}_E : (p_P, q_E) \in \mathcal{I}\}$. The game \mathcal{G} begins by the Pursuer choosing a position $p_P \in \mathcal{I}_P$, and then the Evader choosing a position $q_E \in \mathcal{I}_E(p_P)$.

- (6) After each side has chosen its initial position, the sides move alternately with the Pursuer moving first. Each side, on its turn, must choose an allowed move given the current state of the game.
- (7) The rules of the game specify when the Pursuer has caught the Evader. That is, there is a subset \mathcal{F} of *final positions*. The Pursuer wins \mathcal{G} if, at any time, the current position of the game belongs to \mathcal{F} . The Evader wins the current position never belongs to \mathcal{F} .

We only consider generalized Cops and Robbers games where the set \mathcal{P} of positions is finite. All the games we consider are played over a countable sequence of discrete time-steps or *rounds* which are indexed by natural numbers (including 0).

Analogously to combinatorial games, a generalized Cops and Robbers game can be described as a game played on a bipartite digraph (for example, see [20]). For a given generalized Cops and Robbers game \mathcal{G} , construct the *state digraph*, $\mathcal{D}_{\mathcal{G}}$ as follows. The vertex set V is the disjoint union $\mathcal{S}_P \cup \mathcal{S}_E$, where \mathcal{S}_P is the set of states of the game when it is the Pursuer's turn to move, and \mathcal{S}_E is the set of states of the game when it is the Evader's turn to move. The directed edges of $\mathcal{D}_{\mathcal{G}}$ have one end in \mathcal{S}_P and the other in \mathcal{S}_E ; there is a directed edge from s to t whenever the rules of the game allow a move so that the state of the game changes from s to t . After the game has started, we can imagine the state of the game being represented by a unique token placed on a vertex of $\mathcal{D}_{\mathcal{G}}$. Each player, on his turn, makes a move *from* the current vertex s *to* another vertex t by sliding the token along a directed edge from s to t . The Pursuer wins if and only if the token is eventually on a vertex $((p_P, q_E), X)$ with $(p_P, q_E) \in \mathcal{F}$. Otherwise, the Evader wins.

All generalized Cops and Robbers game are thus, in some sense, games played on directed graphs. As with combinatorial games [20, 21], a labelling of the state digraph can be used to determine if the Pursuer has a winning strategy (also see [2, 15]). We describe the labelling procedure in the case of generalized Cops and Robbers games. Initially, the vertices $((p_P, q_E), X)$ with $(p_P, q_E) \in \mathcal{F}$ have $label(X) = 0$, and all other vertices X have $label(X) = \infty$. We use the notation $N^+(u)$ for the set of out-neighbours of a vertex u . The following process is then repeated until no labels change:

- (1) If $X \in \mathcal{S}_P$, then set $label(X) = 1 + \min_{Y \in N^+(X)} label(Y)$.
- (2) If $X \in \mathcal{S}_E$, then set $label(X) = 1 + \max_{Y \in N^+(X)} label(Y)$.

The intuition behind the labels is that the Pursuer's optimal strategy is to move so that the game is over as quickly as possible, and the Evader's optimal strategy is to move so the game lasts as long as possible. It is straightforward to prove by induction that the final labels are the number of moves that the game will last from a given position, assuming both sides play optimally (see Theorem 3). The Pursuer has a winning strategy if and only if he can choose a starting position $p_P \in \mathcal{I}_P$ so that $label(P) < \infty$ for all $q_E \in \mathcal{I}_E(p_P)$.

We next provide some examples of generalized Cops and Robbers games. The list is by no means exhaustive.

- (1) *Cops and Robbers* [1, 16, 17, 18, 19]. Players moving to neighbouring vertices, and the final states represent any move of a cop onto the vertex containing the robber.
- (2) *Distance- k Cops and Robbers* [3, 6]. In this variation, the cops and robber move as in the original game, but the final states include the case when the robber is within distance k of some cop, where k is a fixed positive integer.

- (3) *Tandem-win Cops and Robbers* [9, 11]. In this variation, there are pairs of cops who must remain within distance one of each other. This alters only the set of allowed positions of the cops.
- (4) *Cops and Robbers with traps* [10]. Some devices are deployed by the cops so that if the robber enters a node with a “trap” he loses the game. Here, we restrict the allowed moves of the robber. In particular, if T is the set of vertices with traps, then \mathcal{F} contains $V_P \times T$ (along with the usual final states in Cops and Robbers).
- (5) *Eternal domination* [13]. Fix a graph G . The positions for the Pursuer are the vertices of G . The positions for the Evader are the k -subsets of $V(G)$, where $k \geq 1$ is an integer. The vertices in these k -subsets are regarded as each holding a guard. The positions for the Pursuer are regarded as vertices where the Evader must locate a guard on his next move. The allowed starting positions are the pairs (x, X) with $x \in X$. On his turn, the Pursuer can move to any vertex from any other vertex. On his turn, the Evader can move from position X_1 to position Y_1 when every guard in X_1 can slide along an edge (maybe a loop) so that the resulting configuration is Y_1 (or whatever the variant of the game dictates; for example, only one guard may move). The final states are the pairs (x, Y) with $x \notin Y$. The graph G has an *eternal dominating set of cardinality k* if and only if the Evader has a winning strategy.
- (6) *Seepage* [4, 7]. This game is played on a directed acyclic graph, with a single source (vertex of in-degree zero) and a fixed number of sinks (vertices of out-degree zero). the Evader is called *sludge* and there is some number of *greens* (the Pursuer). The sludge begins at the source and, on each turn, moves along directed edges to any out-neighbours not protected by the greens, or simply does not move. On their turn, the greens protect some vertices not occupied by sludge by moving to them, if possible, (one vertex per green; not necessarily an adjacent vertex); once a vertex is protected, it remains that way to the end of the game. The final positions are those where the greens have prevented sludge from entering a sink.

3. A RELATIONAL CHARACTERIZATION

We now give a relational characterization which generalizes the one given in [17] for Cops and Robbers in the case of one cop, and later in [8] for more than one cop. We emphasize that the characterization we provide applies to all generalized Cops and Robbers games as defined in Section 2. We note that Cops and Robbers is a kind of position independent game where the relational characterization takes on specific additional properties; see Section 4.

Let \mathcal{G} be a generalized Cops and Robbers game. Similarly to [8, 17], define a non-decreasing sequence of relations from \mathcal{P}_E to \mathcal{P}_P , indexed by natural numbers, as follows:

- (1) $q_E \preceq_0 p_P$ if and only if $(p_P, q_E) \in \mathcal{F}$.
- (2) Suppose $\preceq_0, \preceq_1, \dots, \preceq_{i-1}$ have all been defined for some $i \geq 1$. Define $q_E \preceq_i p_P$ if $(p_P, q_E) \in \mathcal{F}$, or if $((p_P, q_E), R) \in \mathcal{S}$ and, for every $x_E \in \mathcal{A}_E(p_P, q_E)$ either $(p_P, x_E) \in \mathcal{F}$ or there exists $w_P \in \mathcal{A}_P(p_P, x_E)$ such that $x_E \preceq_j w_P$ for some $j < i$.

By definition \preceq_i contains \preceq_{i-1} for each $i \geq 1$. Since $\mathcal{P}_E \times \mathcal{P}_P$ is finite, there exists t such that $\preceq_t = \preceq_k$ for all $k \geq t$. Define $\preceq = \preceq_t$.

We next use this sequence of relations to derive a method of determining the winner of a given generalized Cops and Robbers game. For $i > 0$, the intuition behind the definition of \preceq_i can be phrased as: *if the game is not over, for every move that the Evader can make*

from this position, either the game ends or the Pursuer has a response that leads to a win after he has moved j more times. Since the Pursuer is next to move once a start position $(p_P, q_E) \in \mathcal{I}$ has been determined, it is not necessary that $q_E \preceq p_P$ in order for the Pursuer to have a winning strategy. Instead he needs to be able to choose his initial position p_P in such a way that no matter which initial position is chosen by the Evader, he has a move to a position $w_P \in \mathcal{A}_P(p_P, q_E)$ such that $q_E \preceq w_P$.

Theorem 1. *The Pursuer has a winning strategy in the generalized Cops and Robbers game \mathcal{G} if and only if there exists $p_P \in \mathcal{I}_P$ such that, for all $q_E \in \mathcal{I}_E(p_P)$, either $(p_P, q_E) \in \mathcal{F}$ or there exists $w_P \in \mathcal{A}_P(p_P, q_E)$ such that $q_E \preceq w_P$.*

Proof. We prove the contrapositive of the forward implication. Suppose that the condition does not hold. Then for all $p_P \in \mathcal{I}_P$, there exists $q_E \in \mathcal{I}_E(p_P)$ such that there is no $w_P \in \mathcal{A}_P(p_P, q_E)$ with $q_E \preceq w_P$. If the Evader chooses such a position then, no matter the position v_P to which the Pursuer moves, the state of the game is $((v_P, q_E), R)$ with $q_E \not\preceq v_P$. By definition of the sequence of relations, the Evader can move to $x_E \in \mathcal{A}_E(v_P, q_E)$ for which there is no $y_P \in \mathcal{A}_P(v_P, x_E)$ such that $x_E \preceq y_P$. By induction, it never occurs that the Pursuer is on p_P and the Evader is on q_E such that $q_E \preceq_0 p_P$; hence, the Evader has a winning strategy.

For the reverse implication, we first prove by induction that if it is the Evader's move and the state of the game is (p_P, q_E) with $q_E \preceq_i p_P$, then the Pursuer wins after making i more moves. This is clear when $i = 0$. Assume the statement holds for all natural numbers j with $0 \leq j \leq i - 1$. By definition of the sequence of relations, since $q_P \preceq_i p_E$ and $q_P \not\preceq_{i-1} p_E$, the Evader has a move to a position $w_E \in \mathcal{A}_E(p_E, q_P)$ such that, for all positions $x_P \in \mathcal{A}(p_P, w_E)$ to which the Pursuer can subsequently move, $i - 1 = \min\{j : w_E \preceq_j x_P\}$. Hence, by definition of \preceq_i , the statement holds for i .

Suppose there exists $p_P \in \mathcal{I}_P$ such that, for all $q_E \in \mathcal{I}_E(p_P)$, either $(p_P, q_E) \in \mathcal{F}$ or there exists $w_P \in \mathcal{A}_P(p_P, q_E)$ such that $q_E \preceq w_P$. After the start state has been determined it is the Pursuer's move. The condition guarantees that if the game is not over, then it is always possible for the Pursuer to move to a state from which he has a winning strategy. Hence, the reverse implication holds. \square

The definition of \preceq is similar to the labelling procedure described in Section 2, except that the indices in the sequence take into account a move by each side. Assuming that the Pursuer has a winning strategy, the sequence of relations can be used to determine the length of the game in terms of the number of moves that he must make in order to win assuming both sides play optimally. For an allowed start position $(p_P, q_E) \in \mathcal{I}$, define

$$\ell(p_P, q_E) = \begin{cases} \min\{j : q_E \preceq_j p_P\} & \text{if } q_E \preceq_j p_P, \\ 1 + \max_{w_P \in \mathcal{A}_P(p_P, q_E)} \min\{j : q_E \preceq_j w_P\} & \text{otherwise.} \end{cases}$$

We have the following corollary.

Corollary 2. *Suppose the Pursuer has a winning strategy in the generalized Cops and Robbers game \mathcal{G} . Assuming optimal play the length of the game is*

$$\min_{p_P \in \mathcal{I}_P} \max_{q_E \in \mathcal{I}_E(p_P)} \ell(p_P, q_E).$$

Proof. As noted in the proof of Theorem 1, by definition of the increasing sequence of relations, if $x \preceq_i y$ and $x \not\preceq_{i-1} y$, then the Evader has a move to a position $z \in \mathcal{A}_E(y, x)$

such that

$$i - 1 = \min_{w \in \mathcal{A}_P(z, x)} \{j : z \preceq_j w\}.$$

Suppose the Pursuer chooses $p_P \in \mathcal{I}_P$, and the Evader subsequently chooses an allowed starting position $q_E \in \mathcal{I}_E(p_P)$. If $q_E \preceq_i p_P$, then assuming optimal play, the process ends in i more moves by the Pursuer. If, on the other hand, $q_E \not\preceq_i p_P$, then assuming optimal play, the Pursuer will move to a position w_P for which the value of j such that $q_E \preceq_j w_P$ is minimized, and the process ends in $j + 1$ more moves by the Pursuer. \square

We now directly establish a connection between the labelling of the state digraph and the sequence of relations.

Theorem 3. *Let \mathcal{G} be a generalized Cops and Robbers game. The label of the vertex $((p_P, q_E), R) \in V(\mathcal{D}_{\mathcal{G}})$ equals k if and only if $q_E \preceq_{\lceil \frac{k}{2} \rceil} p_P$ and $q_E \not\preceq_{\lceil \frac{k}{2} \rceil - 1} p_P$ when $k \geq 1$.*

Proof. The result will follow if we can establish the following items: (i) if $q_E \preceq p_P$, then $((p_P, q_E), R)$ is labelled with a natural number; and (ii) if the label of the vertex $((p_P, q_E), R) \in V(\mathcal{D}_{\mathcal{G}})$ equals k , then $q_E \preceq_{\lceil \frac{k}{2} \rceil} p_P$ and $q_E \not\preceq_{\lceil \frac{k}{2} \rceil - 1} p_P$ when $k \geq 1$.

To prove (i) we argue by induction on n that if $q_E \preceq_n p_P$, then $((p_P, q_E), R)$ is labelled with a natural number. The statement is true when $n = 1$ by the labelling procedure. Suppose, for some $m \geq 1$, that it is true when $n = m - 1$, and that $q_E \preceq_m p_P$.

Then for every $w_E \in \mathcal{A}_E(p_P, q_E)$ either $(p_P, w_E) \in \mathcal{F}$, or there exists $x_P \in \mathcal{A}_P(p_P, w_E)$ such that $w_E \preceq_j x_P$ for some $j < m$. Hence, by the labelling procedure, for every $w_E \in \mathcal{A}_E(p_P, q_E)$ either $\text{label}((p_P, w_E), L) = 0$, or there exists $x_P \in \mathcal{A}_P(p_P, w_E)$ such that (x_P, w_E) is labelled with a natural number. In the latter case, by the labelling procedure again, the vertex $((p_P, w_E), L)$ is labelled with a natural number, since every out-neighbour of the vertex $(p_P, w_E), R)$ is labelled with a natural number. The statement now follows by induction.

The proof of (ii) is by induction on k . It follows from the definitions that the statement is true when $k = 0$. If $k = 1$, then we have $(p_P, q_E) \notin \mathcal{F}$, but for every $w_E \in \mathcal{A}_E(p_P, q_E)$, the position $(p_P, w_E) \in \mathcal{F}$. Hence, by definition of the sequence of relations, $q_E \preceq_1 p_P$. Since $(p_P, q_E) \notin \mathcal{F}$, we have $q_E \not\preceq_0 p_P$. Therefore, the statement is also true when $k = 1$.

Suppose, for some $t \geq 2$, the statement is true when $0 \leq k \leq t - 1$, and that $\text{label}((p_P, q_E), R) = t$. Then by the definition of the labelling,

$$t - 1 \geq \max_{w_E \in \mathcal{A}_E(p_P, q_E)} \text{label}((p_P, w_E), L)$$

and equality is achieved at least once. By definition of the labelling again, for any the Evader position $w_E \in \mathcal{A}_E(p_P, q_E)$,

$$t - 2 \geq \min_{x_P \in \mathcal{A}_P(p_P, w_E)} \text{label}((w_P, w_E), R)$$

and equality is achieved at least once for each w_E with $\text{label}((p_P, w_E), L) = t - 1$. Thus, by the induction hypothesis, for every $w_E \in \mathcal{A}_E(p_P, q_E)$, either $(p_P, w_E) \in \mathcal{F}$, or there exists $x_P \in \mathcal{A}_P(p_P, w_E)$ such that $w_E \preceq_j x_P$ for some $j < t$, and at least one such j equals $t - 1$. Therefore, $q_P \preceq_t p_P$ and $q_E \not\preceq_{t-1} p_P$. The statement now follows by induction. \square

We note that the labels on the vertices in \mathcal{S}_P are uniquely determined by the labelling procedure once the labels on the vertices in \mathcal{S}_E are known.

4. POSITION INDEPENDENT GAMES

In games like Cop and Robbers and many of its variants, each side has the same set of available moves irrespective of the position of the other side (some of these moves may be much better than other, however; for example, a bad but allowed move of the robber is to move the position of a cop).

We define a generalized Cops and Robbers game \mathcal{G} to be *position independent* if for all $p_P \in \mathcal{P}_P$ and $q_E \in \mathcal{P}_E$, whenever $(p_P, q_E) \notin \mathcal{F}$ the set $\mathcal{A}_P(p_P, q_E)$ depends only on p_P and the set $\mathcal{A}_E(p_P, q_E)$ depends only on q_E . That is, if the game is not over, from any position of the game, the set of available moves for a side does not depend on the position of the other side. A game is called *position dependent* if it is not position independent. For example, Cops and Robbers is position independent, while Seepage is position dependent.

In a position independent game, the set of positions of the game can be taken to be $\mathcal{P}_P \times \mathcal{P}_E$. Furthermore, because of position independence one can define sets \mathcal{M}_P and \mathcal{M}_E of *allowed moves* for the Pursuer and the Evader, respectively; for example $(p_P, p'_P) \in \mathcal{M}_P$ if and only if $p'_P \in \mathcal{A}_P(p_P, q_E)$ (for all $q_E \in \mathcal{P}_E$). The ordered pairs $G_P = (P_P, M_P)$ and $G_E = (P_E, M_E)$ are the *position digraphs* for the Pursuer and the Evader, respectively. For all $p_P \in \mathcal{P}_P$ and $q_E \in \mathcal{P}_E$, we have that $\mathcal{A}_P(p_P, q_E) = N_{G_P}^+(p_P)$ and $\mathcal{A}_E(p_P, q_E) = N_{G_E}^+(q_E)$.

For digraphs G and H , the *categorical product*, $G \times H$, has vertices $V(G) \times V(H)$ and a directed edge $(a, b)(c, d)$ whenever $(a, c) \in E(G)$ and $(b, d) \in E(H)$. The categorical product $\mathcal{R}_{\mathcal{G}} = G_P \times G_E$ is the *round summary digraph* for the game \mathcal{G} . The vertices of this digraph represent the state of the game at the end of each round consisting of a move by each side, and assuming the Evader is next to move. The edges represent possible transitions between states in consecutive rounds. Not all of the edges necessarily make sense. For example, it may be that one cannot make the transition from state s_1 to state s_2 without a state in F arising in the middle of the round. Directed edges between final positions also make no sense. Technically, then, the position digraph should be a subgraph of $\mathcal{R}_{\mathcal{G}}$. We will not make this distinction because the extra edges will not affect our treatment.

The following is a consequence of Theorem 1.

Corollary 4. *Let \mathcal{G} be a position independent generalized Cops and Robbers game. If G_P is strongly connected and there exists $X \subseteq P_P$ such that $\mathcal{I} = X \times P_E$, then the Pursuer has a winning strategy in \mathcal{G} if and only if $\preceq = V(\mathcal{R}_{\mathcal{G}}) = P_P \times P_E$.*

Proof. The proof of the reverse direction is analogous to the proof of the forward direction in Theorem 1, and so is omitted. For the forward direction, suppose the Pursuer has a winning strategy. Then there exists an allowed starting position, say p_P , for the Pursuer such that for all $q_E \in P_E$, either $(p_P, q_E) \in \mathcal{F}$ or there exists $w_P \in N_{G_P}^+(p_P)$ and a natural number j such that $q_E \preceq_j w_P$. Therefore, there is a natural number i such that $q_E \preceq_i p_P$ for all $q_E \in P_E$. It now follows by induction that all $q_E \preceq_{i+k} x$ for all $q_E \in P_E$ for all x joined to p_P by a directed path of length k . The result now follows as G_P is strongly connected. \square

The position independent case gives rise to an algorithm which is a polynomial time function of the number of positions, assuming the number of allowed moves of the Pursuer and the Evader is not too large. In the case of the Cops and Robbers game played with k cops on a graph of order n (where k is fixed and not a function of n), Theorem 5 gives a $O(n^{2k+2})$ algorithm for determining if the cops have a winning strategy. This matches the bound on the complexity of the algorithms presented in [3, 15].

Theorem 5. *Let \mathcal{G} be a position independent generalized Cops and Robbers game. Given the graphs G_P and G_E , if $N_{G_P}^+(p_P)$ and $N_{G_E}^+(p_E)$ can be obtained in time $O(f_P(\mathcal{P}_P))$ and $O(f_E(\mathcal{P}_E))$, respectively, then there is a $O(|\mathcal{P}_P| \cdot |\mathcal{P}_E| \cdot f_P(|\mathcal{P}_P|) \cdot f_E(|\mathcal{P}_E|))$ algorithm to determine if the Pursuer has a winning strategy, and the length of the game assuming optimal play.*

Proof. We describe an algorithm to determine if the Pursuer has a winning strategy in a position independent game. Define the matrix $\mathcal{M}_{\mathcal{G}}$, with rows indexed by elements of \mathcal{P}_P and columns indexed by elements of \mathcal{P}_E , to record the sequence of relations. If $(p_P, q_E) \in \mathcal{F}$, then the (p_P, q_E) entry of $\mathcal{M}_{\mathcal{G}}$ is zero. For $i > j$, the (p_P, q_E) entry of $\mathcal{M}_{\mathcal{G}}$ equals i if for every $x_E \in N_{G_E}^+(q_E)$, there exists $y_P \in N_{G_P}^+(p_P)$ such that the (x_E, y_P) entry of $\mathcal{M}_{\mathcal{G}}$ equals j . The matrix $\mathcal{M}_{\mathcal{G}}$ has $|\mathcal{P}_P \times \mathcal{P}_E|$ entries. For $i = 1, 2, \dots$, and until no entries in the matrix change, each entry in $\mathcal{M}_{\mathcal{G}}$ not yet assigned a value must be tested to see if it can be set to i . It can be if, for each $x_E \in N_{G_E}^+(q_E)$, there exists $y_P \in N_{G_P}^+(p_P)$ such that the (x_E, y_P) entry of $\mathcal{M}_{\mathcal{G}}$ equals $j < i$. For each i we can update $\mathcal{M}_{\mathcal{G}}$ row-by-row.

By Corollary 4, the Pursuer has a winning strategy if and only if every entry of $\mathcal{M}_{\mathcal{G}}$ is eventually assigned a value. By definition of the sequence of relations, one can conclude that the Pursuer has a winning strategy as soon as some row of $\mathcal{M}_{\mathcal{G}}$ has a value for each entry. More information can be obtained by filling in the entire matrix. By Corollary 2, the length of the game, assuming optimal play, is $\max_{p_P \in \mathcal{P}_P} \min_{q_E \in \mathcal{P}_E} \mathcal{M}_{\mathcal{G}}(p_P, q_E)$.

We now consider the complexity of this algorithm. It takes $|\mathcal{F}| \leq |\mathcal{P}_P \times \mathcal{P}_E|$ steps to set the entries corresponding to the final positions to zero. For each i , every entry not yet assigned a value must be tested to see if it can be set to i . The procedure attempts to fill in $\mathcal{M}_{\mathcal{G}}$ row-by-row. For each row p_P , suppose it takes $f_P(|\mathcal{P}_P|)$ time to obtain $N_{G_P}^+(p_P)$. Possibly all $|\mathcal{P}_E|$ entries in the row must be tested. To test whether the entry in column q_E can be assigned a value requires obtaining the list of all possible the Evader moves from q_E . Suppose this takes time $f_E(|\mathcal{P}_E|)$. Then processing row p_P takes time at most $O(f_P(|\mathcal{P}_P|) \cdot |\mathcal{P}_E| \cdot f_E(|\mathcal{P}_E|))$. There are $|\mathcal{P}_P|$ rows and possibly as many as $|\mathcal{P}_P \times \mathcal{P}_E|$ values for i . Hence, given the graphs G_P and G_E , the algorithm takes time at most $O(|\mathcal{P}_P| \cdot |\mathcal{P}_E| \cdot f_P(|\mathcal{P}_P|) \cdot f_E(|\mathcal{P}_E|))$, as desired. \square

As an application of Theorem 5, consider the game of Cops and Robber on a reflexive graph G with a fixed positive number k of cops. The graph G_E is the reflexive graph G and the graph G_P is the k -fold categorical product of G with itself. We can take f_P and f_E to be the maximum degree of G_P and G_E , respectively. Hence, f_P is $O(n^k)$ and f_E is $O(n)$. Finally, $|\mathcal{F}| = 2n^2$ and $|\mathcal{P}_E \times \mathcal{P}_P| = n^{k+1}$. Thus, given G_P and G_E , we can decide if the Pursuer has a winning strategy, and the length of the game assuming optimal play, in time

$$O(n^{k+1}(n \cdot n^k)) = O(n^{2k+2}).$$

Note, however, a succinct description of the game would consist only of the graph G on n vertices. The graph G_P would need to be constructed, which can be accomplished in time $O(n^{2k})$. The overall complexity does not change because

$$O(n^{2k} + n^{k+1}(n \cdot n^k)) = O(n^{2k+2}).$$

We can also obtain a vertex elimination ordering for generalized Cops and Robbers games, analogous to cop-win orderings characterizing cop-win graphs; see [17]. In a position independent generalized Cops and Robbers game \mathcal{G} , a vertex (p_P, q_E) of $\mathcal{R}_{\mathcal{G}}$ is *removable with respect to* $X \subseteq V(\mathcal{R}_{\mathcal{G}})$ if either

- (1) it is in \mathcal{F} , or
- (2) for every $x_E \in N_{G_E}^+(q_E)$ either $(p_P, x_E) \in \mathcal{F}$ or there exists $y_P \in N_{G_P}^+(p_P)$ such that (y_P, x_E) is in X .

A *removable vertex ordering* for \mathcal{R}_G is a sequence of vertices in which each element (p_P, q_E) is removable with respect to the set X of vertices that precede it in the sequence, and there exists $p_P \in \mathcal{I}_P$ so that, for all $q_E \in \mathcal{I}_E(p_P)$, either $(p_P, q_E) \in \mathcal{F}$ or there exists $w_P \in \mathcal{A}(p_P, q_E)$ such that (w_P, q_E) belongs to the sequence.

If the Pursuer has a winning strategy, then listing the vertices of \mathcal{S}_G so that the pairs in \preceq_0 are followed by those in $\preceq_1 \setminus \preceq_0$, then those in $\preceq_2 \setminus \preceq_1$, and so on, is a removable vertex ordering for \mathcal{S}_G . Other orders may be possible; for example, it may be that not all pairs in \preceq_0 need to be listed before a pair in \preceq_1 can be listed.

We now have the following vertex-ordering characterization of games where the Pursuer has a winning strategy.

Corollary 6. *The Pursuer has a winning strategy in the position independent generalized Cops and Robbers game \mathcal{G} if and only if \mathcal{R}_G admits a removable vertex ordering.*

Proof. Suppose that the Pursuer has a winning strategy. Then by Theorem 1, there exists $p_P \in \mathcal{I}_P$ such that for all $q_E \in \mathcal{I}_E(p_P)$ either $(p_P, q_E) \in \mathcal{F}$, or there exists $w_P \in \mathcal{A}(p_P, q_E)$ with $q_E \preceq w_P$. Listing the vertices of \mathcal{S}_G so that the pairs in \preceq_0 are followed by those in $\preceq_1 \setminus \preceq_0$, then those in $\preceq_2 \setminus \preceq_1$, and so on, is a removable vertex ordering for \mathcal{R}_G .

On the other hand, suppose \mathcal{R}_G admits a removable vertex ordering. Then there is a sequence vertices such that each element (p_P, q_E) is removable with respect to the set X of vertices that precede it in the sequence, and there exists $p_P \in \mathcal{I}_P$ so that for all $q_E \in \mathcal{I}_E(p_P)$ either $(p_P, q_E) \in \mathcal{F}$ or there exists $w_P \in N_{G_P}^+(p_P)$ such that (w_P, q_E) belongs to the sequence. the Pursuer chooses p_P for his starting position. No matter which position q_E the Evader chooses, if the game is not over then the Pursuer has a move to a position w_P so that (w_P, q_E) belongs to the sequence. By definition of a removable vertex ordering, the position each subsequent move by the Pursuer is closer to the beginning of the sequence. Hence, eventually a position in \mathcal{F} is encountered. \square

5. k -PURSUER GAMES

In this final section, we consider an analogue of the cop number for generalized Cops and Robbers games. First, we need a few definitions. Given a digraph G with subdigraph H , a *retraction onto H* is a mapping $f : G \rightarrow H$ such that f is the identity on H . Let G_1 and G_2 be digraphs, with H_i a subdigraph of G_i for $i = 1, 2$. If there are retractions $f_1 : G_1 \rightarrow H_1$ and $f_2 : G_2 \rightarrow H_2$, then define the product function $g_1 \times g_2 : G_1 \times H_1 \rightarrow G_2 \times H_2$ defined by $(f_1 \times f_2)(x, y) = (f_1(x), f_2(y))$. It is straightforward to see that f_1 and f_2 are retractions, then so is $f_1 \times f_2$.

Let \mathcal{G} be a position independent discrete-time pursuit game. Suppose that G'_P and G'_E are subgraphs of G_P and G_E . We define the *restriction of \mathcal{G} to G'_P and G'_E* to be the discrete-time pursuit game \mathcal{G}' with position digraphs G'_P and G'_E , start states $\mathcal{I}' = \mathcal{I} \cap V(G'_P \times G'_E)$ and final states $\mathcal{F}' = \mathcal{F} \cap V(G'_P \times G'_E)$.

Now suppose that G'_P and G'_E are *induced* subgraphs of G_P and G_E . A *retraction of \mathcal{G} to \mathcal{G}'* is a pair of retractions $\ell : G_P \rightarrow G'_P$ and $r : G_E \rightarrow G'_E$ such that

- (1) If $(x, y) \in \mathcal{I}$ then $(\ell \times r)(x, y) \in \mathcal{I}'$;
- (2) If $(x, y) \in \mathcal{F}$, then $(\ell \times r)(x, y) \in \mathcal{F}'$;

(3) If $(x', y') \in \mathcal{F}'$ then $(\ell \times r)^{-1}(x', y') \subseteq \mathcal{F}$.

If there is a retraction of \mathcal{G} to \mathcal{G}' , then \mathcal{G}' is a *retract* of \mathcal{G} .

Lemma 7. *If \mathcal{G}' is a retract of \mathcal{G} and the Evader has a winning strategy in \mathcal{G}' , then the Evader has a winning strategy in \mathcal{G} .*

Proof. The Evader plays the game \mathcal{G} as if playing \mathcal{G}' , regarding the Pursuer's position p_P as $\ell(p_P)$ (thinking of this as the Pursuer's "shadow"). Since the Evader has a winning strategy, a state in \mathcal{F}' is never encountered. By definition of retraction, a position in \mathcal{F} is never encountered. \square

The following corollary follows immediately from Lemma 7.

Corollary 8. *If the Pursuer has a winning strategy in \mathcal{G} , then the Pursuer has a winning strategy in every retract of \mathcal{G} .*

As an example, consider a game, \mathcal{G} , Cops and Robbers, with one cop, played on a finite graph G . Suppose that H is a given subgraph of G and there is a retraction r of G onto H .

We now provide an analogue of the cop number. Given a pursuit game \mathcal{G} and an integer $k \geq 1$, the k -pursuer game \mathcal{G}_k is defined as follows.

- (1) The set of allowed positions for the Pursuer is the k -fold product \mathcal{P}_P^k .
- (2) The set of allowed positions for the Evader is \mathcal{P}_E .
- (3) The set of start positions is $\mathcal{I}_k = \{(p_1, p_2, \dots, p_k, q) : (p_i, q) \in \mathcal{I}, 1 \leq i \leq k\}$.
- (4) The set of final positions is $\mathcal{F}_k = \{(p_1, p_2, \dots, p_k, q) : (p_i, q) \in \mathcal{F}, \text{ for some } i \in \{1, 2, \dots, k\}\}$.
- (5) the Pursuer can move so that the position changes from $(p_1, p_2, \dots, p_k, q)$ to $(w_1, w_2, \dots, w_k, q)$ whenever, for $1 \leq i \leq k$, the move so that the position changes from (p_i, q) to (w_i, q) is allowed in \mathcal{G} .
- (6) the Evader can move so that the position changes from $(p_1, p_2, \dots, p_k, q)$ to $(p_1, p_2, \dots, p_k, x)$ whenever, for $1 \leq i \leq k$, the move so that the position changes from (p_i, q) to (p_i, x) is allowed in \mathcal{G} .

The following theorem was proved in [12].

Theorem 9. *Computing $p(\mathcal{G})$ is **NP**-hard for finite Cops and Robber games \mathcal{G} .*

Since Cops and Robber is a pursuit game, the following corollary is immediate.

Corollary 10. *Computing $p(\mathcal{G})$ is **NP**-hard for finite pursuit games \mathcal{G} .*

We stress that this result only states that, unless $\mathbf{P} = \mathbf{NP}$, there is no general algorithm that computes $p(\mathcal{G})$ for all pursuit games \mathcal{G} . It makes no statement about the existence of such an algorithm for particular collections of games.

REFERENCES

- [1] M. Aigner, M. Fromme, A game of cops and robbers, *Discrete Applied Mathematics* **8** (1984) 1–12.
- [2] A. Berarducci, B. Intriglia, On the cop number of a graph, *Advances in Applied Mathematics* **14** (1993) 389–403.
- [3] A. Bonato, E. Chiniforooshan, P. Prałat, Cops and Robbers from a distance, *Theoretical Computer Science* **411** (2010) 3834–3844.
- [4] A. Bonato, D. Mitsche, P. Prałat, Vertex-pursuit in random directed acyclic graphs, *SIAM Journal on Discrete Mathematics* **27** (2013) 732–756.

- [5] A. Bonato, R.J. Nowakowski, *The Game of Cops and Robbers on Graphs*, American Mathematical Society, Providence, Rhode Island, 2011.
- [6] J. Chalopin, V. Chepoi, N. Nisse, Y. Vaxès, Cop and robber games when the robber can hide and ride, *SIAM Journal on Discrete Mathematics* **25** (2011) 333-359.
- [7] N.E. Clarke, S. Finbow, S.L. Fitzpatrick, M.E. Messinger, R.J. Nowakowski, Seepage in directed acyclic graphs, *Australasian Journal of Combinatorics* (2009) **43** 91–102.
- [8] N.E. Clarke, G. MacGillivray, Characterizations of k -copwin graphs, *Discrete Mathematics* **312** (2012) 1421-1425.
- [9] N.E. Clarke, R.J. Nowakowski, A tandem version of the Cops and Robber game played on products of graphs, *Discussiones Mathematicae Graph Theory* **25** (2005) 241–249.
- [10] N.E. Clarke, R.J. Nowakowski, Cops, robber, and traps, *Utilitas Mathematica* **60** (2001) 91–98.
- [11] N.E. Clarke, R.J. Nowakowski, Tandem-win Graphs, *Discrete Mathematics* **299** (2005) 56–64.
- [12] F.V. Fomin, P.A. Golovach, J. Kratochvíl, N. Nisse, K. Suchan, Pursuing fast robber in graphs, *Theoretical Computer Science* **411** (2010) 1167–1181.
- [13] W. Goddard, S.M. Hedetniemi, S.T. Hedetniemi, Eternal security in graphs, *Journal of Combinatorial Mathematics and Combinatorial Computing* **52** (2005) 169–180.
- [14] A.S. Goldstein, E. M. Reingold, The complexity of pursuit on a graph, *Theoretical Computer Science* **143** (1995) 93–112.
- [15] G. Hahn, G. MacGillivray, A note on k -cop, ℓ -robber games on graphs, *Discrete Mathematics* **306** (2006) 2492–2497.
- [16] S. Neufeld, R.J. Nowakowski, A vertex-to-vertex pursuit game played with disjoint sets of edges, *Finite and infinite combinatorics in sets and logic* (Banff, AB, 1991), Kluwer Acad. Publ., 1993, 411, 299-312
- [17] R.J. Nowakowski, P. Winkler, Vertex to Vertex Pursuit in a Graph, *Discrete Mathematics* **43** (1983) 23–29.
- [18] A. Quilliot, Jeux et pointes fixes sur les graphes, *Thèse de 3ème cycle*, Université de Paris VI, 1978, 131–145.
- [19] A. Quilliot, *Problèmes de jeux, de point Fixe, de connectivité et de représentation sur des graphes, des ensembles ordonnés et des hypergraphes*, Thèse d'Etat, Université de Paris VI, 1983, 131–145.
- [20] C.A.B. Smith, Graphs and composite games, *Journal of Combinatorial Theory* **1** (1966) 51–81.
- [21] H. Steinhaus, Definitions for a theory of games and pursuit, *Myśl Akad.Lwów* **1** (1925) 13–14; reprinted in *Naval Research Logistics Quarterly* **7** (1960) 105–108.

DEPARTMENT OF MATHEMATICS, RYERSON UNIVERSITY, TORONTO, ON, CANADA
E-mail address: abonato@ryerson.ca

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF VICTORIA, VICTORIA, BC, CANADA
E-mail address: gmacgill@uvic.ca