

Available online at www.sciencedirect.com



Nonlinear Analysis 65 (2006) 1425-1439



www.elsevier.com/locate/na

Central configurations and total collisions for quasihomogeneous *n*-body problems

Florin Diacu^{a,*}, Ernesto Pérez-Chavela^b, Manuele Santoprete^c

^a Department of Mathematics and Statistics, University of Victoria, Victoria, Canada ^b Departamento de Matemáticas, UAM–Iztapalapa, A.P. 55–534, 09340 Iztapalapa, Mexico, D.F., Mexico ^c Department of Mathematics, University of California, Irvine, USA

Received 10 August 2005; accepted 17 October 2005

Abstract

We consider *n*-body problems given by potentials of the form $\frac{\alpha}{r^a} + \frac{\beta}{r^b}$ with *a*, *b*, α , β constants, $0 \le a < b$. To analyze the dynamics of the problem, we first prove some properties related to central configurations, including a generalization of Moulton's theorem. Then we obtain several qualitative properties for collision and near-collision orbits in the Manev-type case a = 1. At the end we point out some new relationships between central configurations, relative equilibria, and homothetic solutions. (© 2005 Elsevier Ltd. All rights reserved.

MSC: 70F10; 70H05

Keywords: Central configuration; Relative equilibria; Total collision manifold

1. Introduction

The *n*-body problem studied here is given by a potential of the form $\frac{\alpha}{r^a} + \frac{\beta}{r^b}$, where *r* is the distance between bodies and *a*, *b*, α , β are constants, $0 \le a < b$ (see [4,12]). In the first part of the paper we treat the general problem, and in the second part we focus on the case a = 1. The function $\frac{\alpha}{r^a} + \frac{\beta}{r^b}$, called quasihomogeneous because of being the sum of homogeneous functions of different degrees, generalizes classical potentials, such as those of

^{*} Corresponding author. Tel.: +1 604 721 7437; fax: +1 604 721 8962.

E-mail addresses: diacu@math.uvic.ca (F. Diacu), epc@xanum.uam.mx (E. Pérez-Chavela), msantopr@math.uci.edu (M. Santoprete).

⁰³⁶²⁻⁵⁴⁶X/\$ - see front matter © 2005 Elsevier Ltd. All rights reserved. doi:10.1016/j.na.2005.10.023

Newton, Coulomb, Birkhoff, Manev, Van der Waals, Libhoff, Schwarzschild, and Lennard-Jones. Thus, the applicability of the quasihomogeneous *n*-body problem ranges from celestial mechanics and atomic physics to chemistry and crystallography.

Although many properties of the Newtonian n-body problem have a correspondent in the homogeneous case, this is not true for nonhomogeneous potentials. On the one hand, the transposition of known results is far from trivial; on the other, new properties show up.

An intriguing aspect we will point out in this paper refers to central configurations, which are crucial for understanding the dynamics of the *n*-body problem (see [13]). The central configurations of the quasihomogeneous potential are in a certain relationship with the central configurations of the homogeneous functions that form this potential. Thus, we will introduce here the notion of simultaneous central configuration and will investigate its connection with the classical concept.

In Section 2, we define the quasihomogeneous *n*-body problem and write down the equations of motion. In Section 3, we introduce the concepts of central and simultaneous central configuration, the latter being specific to quasihomogeneous potentials. Section 4 deals with collinear central configurations. Using critical point theory, we prove a generalization of Moulton's theorem by showing that the number of collinear central configurations of n bodies is n!/2. Starting with Section 5, we restrict our study to Manev-type problems, [5], i.e. those given by potentials of the form $\frac{\alpha}{r} + \frac{\beta}{r^b}$, and show that there are exactly two planar central configurations in the three-body case. Section 6 introduces a framework for the study of collision and near-collision orbits, which is performed in Sections 7 and 8. We study in detail the network of collision solutions and determine the relationship between central configurations, on the one hand, and relative equilibria and homothetic orbits, on the other hand. It is important to note that if in the homogeneous case the correspondence between central configurations and homothetic solutions is one-to-one, this fails to be the case in the quasihomogeneous problem. The relationship between central configurations and relative equilibria remains unchanged, i.e. one-to-one, in the quasihomogeneous case. For Manev-type potentials, homothetic orbits are less likely than in the Newtonian case, in the sense that they show up only for simultaneous central configurations.

2. The quasihomogeneous *n*-body problem

We will start with defining the planar quasihomogeneous *n*-body problem. Consider the linear space

$$\Omega = \left\{ \mathbf{r} = (\mathbf{r}_1, \dots, \mathbf{r}_n) \in (\mathbb{R}^2)^n \middle| \sum_{i=1}^n m_i \mathbf{r}_i = 0 \right\},\tag{1}$$

where $m_i > 0, i = 1, 2, ..., n$, are the masses of the *n* bodies and $\mathbf{r}_i, i = 1, 2, ..., n$, represent their coordinates. Notice that $\sum_{i=1}^{n} m_i \mathbf{r}_i = 0$ fixes the centre of mass at the origin of the coordinate system. Let

$$\Delta_{ij} = \{ (\mathbf{r}_1, \dots, \mathbf{r}_n) \in \Omega \mid \mathbf{r}_i = \mathbf{r}_j \}; \qquad \Delta = \bigcup_{i,j} \Delta_{ij}.$$
(2)

We call Δ the collision set. The potential U of the system is a function defined on the configuration space $\tilde{\Omega} = \Omega \setminus \Delta$ and is given by

$$U = W + V,$$

where W is a homogeneous function of degree $-a, a \ge 0$,

$$W(\mathbf{r}_1,\ldots,\mathbf{r}_n) = \sum_{i< j} \frac{m_i m_j}{\|\mathbf{r}_i - \mathbf{r}_j\|^a},$$
(3)

and V is a homogeneous function of degree -b, b > a,

$$V(\mathbf{r}_1,\ldots,\mathbf{r}_n) = \sum_{i< j} \frac{m_i m_j}{\|\mathbf{r}_i - \mathbf{r}_j\|^b}.$$
(4)

The equations of motion of the *n* bodies define a vector field X on the tangent bundle $T(\tilde{\Omega})$. The configuration space of the system is $\tilde{\Omega}$ and the cotangent bundle is $T^*(\tilde{\Omega})$. Let $\mathbf{p} = M^{-1}\dot{\mathbf{r}}$ be the linear momentum of the system of particles, where M is the diagonal matrix $M = \text{diag}(m_1, m_1, m_2, m_2, \dots, m_n, m_n)$. Then the equations of motion can be written as a Hamiltonian system,

$$\dot{\mathbf{r}} = \frac{\partial H}{\partial \mathbf{p}}$$

$$\dot{\mathbf{p}} = -\frac{\partial H}{\partial \mathbf{r}},$$
(5)

where $H: T^*(\tilde{\Omega}) \to \mathbb{R}$ is the Hamiltonian function given by

$$H(\mathbf{r}, \mathbf{p}) = \frac{1}{2} \mathbf{p}^{t} M^{-1} \mathbf{p} - U(\mathbf{r}).$$
(6)

Here $T = \frac{1}{2}\mathbf{p}^t M^{-1}\mathbf{p}$ is the kinetic energy. The total energy H is a first integral for the system (5); this means that T - U = h (constant) along any orbit. Other integrals are given by the linear momentum, $\sum_{i=1}^{n} m_i \dot{\mathbf{r}}_i$, and by the angular momentum, $J : T \to \mathbf{R}$, defined as

$$J(\mathbf{r}, \mathbf{v}) = \sum_{i=1}^{n} m_i \mathbf{r}_i \times \mathbf{v}_i.$$
(7)

Notice that the relationships for the centre of mass, $\sum_{i=1}^{n} m_i \mathbf{r_i} = 0$, and linear momentum, $\sum_{i=1}^{n} m_i \dot{\mathbf{r}}_i = 0$, together with the energy integral, T - U = h, reduce the dimension of the Hamiltonian system (5) from 4n to 4n - 5. We also introduce the scalar product,

$$\langle \mathbf{r}, \tilde{\mathbf{r}} \rangle = \mathbf{r}^t M \tilde{\mathbf{r}},$$
(8)

which allows us to write the moment of inertia as

$$I = \langle \mathbf{r}, \mathbf{r} \rangle = \sum_{i=1}^{n} m_i \|\mathbf{r}_i\|^2.$$
⁽⁹⁾

3. Central configurations

Central configurations play a crucial role for understanding the dynamics of *n*-body problems [13]. In particular, they have led to important theoretical investigations, such as Saari's conjecture, which has remained open for more than three and a half decades [6], and are connected to Smale's 6th problem [15], originally proposed by Wintner in 1941 [16] (see also [9, 11]). In this section we will define central configurations and analyze the particular aspects this concept encounters in the quasihomogeneous case.

Definition 1. A configuration $\mathbf{r} \in \tilde{\Omega}$ is called central if there is a constant σ such that

$$\nabla U(\mathbf{r}) = \sigma \nabla I(\mathbf{r}). \tag{10}$$

Using the fact that the functions W and V are homogeneous of degree -a and -b, respectively, and applying Euler's theorem for homogeneous functions, we find that

$$\sigma = \frac{-aW(\mathbf{r}) - bV(\mathbf{r})}{2I(\mathbf{r})}.$$
(11)

Definition 2. We call $\mathbf{r} \in \tilde{\Omega}$ a simultaneous central configuration for the potentials *W* and *V* if there are constants σ_1 and σ_2 such that

$$\nabla W(\mathbf{r}) = \sigma_1 \nabla I(\mathbf{r})$$
 and $\nabla V(\mathbf{r}) = \sigma_2 \nabla I(\mathbf{r}).$

Using the fact that W and V are homogeneous functions of degree -a and -b, respectively, we find that

$$\sigma_1 = \frac{-aW(\mathbf{r})}{2I(\mathbf{r})}$$
 and $\sigma_2 = \frac{-bV(\mathbf{r})}{2I(\mathbf{r})}$. (12)

Note that if **r** is a simultaneous central configuration for W and V, then **r** is also a central configuration for U = V + W. The converse is not necessarily true.

Let

$$S_{I_0} = \{ \mathbf{r} \in \Omega \mid \langle \mathbf{r}, \mathbf{r} \rangle = I_0 \}$$

be the sphere relative to the metric given by the scalar product, and denote by

$$S_{I_0}^* = S_{I_0} \setminus \varDelta = \{\mathbf{r} \in \Omega \mid \langle \mathbf{r}, \mathbf{r} \rangle = I_0\}$$

this sphere minus the collision set. Then the central configurations with moment of inertia I_0 can also be defined as the critical points of U_{S_I} , where $U_{S_I} : S_I^* \to \mathbb{R}$ is the restriction of the potential U to $S_{I_0}^*$. Denote by C_n the set of central configurations of the quasihomogeneous *n*-body problem.

Definition 3. We say that two relative equilibria in $S_{I_0}^*$ are equivalent (and belong to the same equivalence class) if they can be made congruent by the induced S^1 action on $S_{I_0}^*$, that is, if one is obtained from the other by a rotation.

Let \tilde{C}_n denote the set of equivalence classes of central configurations. Note that this definition differs from the one used in the Newtonian case (see [1,14]), where two central configurations are called equivalent when one can be obtained from the other by a rotation and/or a homothety. This change is necessary in the quasihomogeneous case because the set C_n is invariant under the action of the group S^1 , but not necessarily under the action of homotheties (see Section 7).

Clearly, I and Δ are invariant under the action of S^1 . Thus, we can conclude that $S_{I_0}^*$ is diffeomorphic to the (2n - 3)-dimensional sphere S^{2n-3} (which is actually an ellipsoid E^{2n-3}) with all the points Δ removed, that is,

$$S_{I_0}^* = E^{2n-3} \setminus (E^{2n-3} \cap \Delta) \approx S^{2n-3} \setminus (S^{2n-3} \cap \Delta).$$

1428

Since U_{S_I} is invariant under the action of S^1 , it defines a map $\tilde{U}_{S_I} : S_{I_0}^*/S^1 \to \mathbb{R}$. If we let $\pi_n : S_{I_0}^* \to S_{I_0}^*/S^1$ denote the canonical projection, $\tilde{\Delta} = \pi(E^{2n-3} \cap \Delta)$, and recalling that $E^{2n-3}/S^1 \approx S^{2n-3}/S^1 \approx \mathbb{C}P^{n-2}$ (the complex projective space), we are led to investigate the critical points of $\tilde{U}_{S_I} : \mathbb{C}P^{n-2} \setminus \tilde{\Delta} \to \mathbb{R}$.

Consequently we can show that the set of equivalence classes of central configurations with fixed moment of inertia I_0 is given by the set of critical points of the map $\tilde{U}_{S_I} : \mathbb{C}P^{n-2} \setminus \tilde{\Delta} \to \mathbb{R}$. More precisely, we have proved the following property:

Proposition 1. For any choice of masses in the planar n-body problem with a quasihomogeneous potential, $n \ge 2$, the set of equivalence classes of central configurations with moment of inertia I_0 is diffeomorphic with the set of critical points of the map $\tilde{U}_{S_I} : \mathbb{C}P^{n-2} \setminus \tilde{\Delta} \to \mathbb{R}$.

4. Moulton's theorem for quasihomogeneous potentials

We will now study collinear central configurations and, using critical point theory, will calculate the number of classes of such configurations for any number n of bodies. The goal of this section is to prove the following result, which generalizes a theorem obtained by Forest Ray Moulton in 1910, [10].

Theorem 1. For any choice of masses in the n-body problem with a quasihomogeneous potential, U, and any given moment of inertia, I_0 , there are exactly n!/2 classes of collinear central configurations. In other words, there are n!/2 classes of central configurations $\mathbf{r} = (\mathbf{r}_1, \ldots, \mathbf{r}_n)$, where all \mathbf{r}_i belong to the same straight line through the origin.

In preparation for the proof, choose some line l in \mathbb{R}^2 . This defines a subset $\Omega_l \subset \Omega$ of $\mathbf{r} = (\mathbf{r}_1, \ldots, \mathbf{r}_n)$ such that each \mathbf{r}_i is on the line l. Let $S_l = S_{I_0} \cap \Omega_l$ and $S_l^* = S_l \setminus \cap (S_l \cap \Delta)$. When S^1 acts on S_{I_0} , only the rotation by π radians leaves S_l invariant. Thus the group \mathbb{Z}_2 acts

on S_l , and on the quotient we have $\mathbb{R}P^{n-2} \setminus \tilde{\Delta} \subset \mathbb{C}P^{n-2} \setminus \tilde{\Delta} \xrightarrow{\tilde{U}_{S_l}} \mathbb{R}$, where $\mathbb{R}P^{n-2} = S_l \setminus \mathbb{Z}_2$ is the real projective space, naturally contained in $\mathbb{C}P^{n-2}$. Here \tilde{U}_{S_l} is induced by the potential energy. From these considerations we obtain:

Lemma 1. The set of equivalence classes of collinear central configurations with moment of inertia I_0 is diffeomorphic to the set of critical points of \tilde{U}_{S_I} : $\mathbb{C}P^{n-2} \setminus \tilde{\Delta} \to \mathbb{R}$ that lie in $\mathbb{R}P^{n-2} \setminus \tilde{\Delta} \subset \mathbb{C}P^{n-2} \setminus \tilde{\Delta}$.

So in order to describe the collinear central configurations, it is sufficient to obtain the critical points of the potential that lie in the real projective space. In general, a critical point of a function restricted to a submanifold is not necessarily a critical point of the function on the ambient manifold. However, we have the following result:

Proposition 2. If $\mathbf{r} \in \mathbb{R}P^{n-2} \setminus \tilde{\Delta}$ is a critical point of $\tilde{U}_{S_I} : \mathbb{R}P^{n-2} \setminus \tilde{\Delta} \to \mathbb{R}$, then \mathbf{r} is also a critical point of $\tilde{U}_{S_I} : \mathbb{C}P^{n-2} \setminus \tilde{\Delta} \to \mathbb{R}$.

To prove this, we first need to know the derivatives of the potential function, which are given below.

Lemma 2. For given masses m_1, \ldots, m_n and U = W + V,

(1) The first derivative of $U: \tilde{\Omega} \to \mathbb{R}$ is

$$DU(\mathbf{r})(\mathbf{v}) = -a \sum_{i \neq j} \frac{m_i m_j}{\|\mathbf{r}_i - \mathbf{r}_j\|^{a+2}} (\mathbf{r}_i - \mathbf{r}_j, \mathbf{v}_i - \mathbf{v}_j)$$
$$-b \sum_{i \neq j} \frac{m_i m_j}{\|\mathbf{r}_i - \mathbf{r}_j\|^{b+2}} (\mathbf{r}_i - \mathbf{r}_j, \mathbf{v}_i - \mathbf{v}_j)$$

for $\mathbf{v} \in \Omega$.

(2) The second derivative is

$$D^{2}U(\mathbf{r})(\mathbf{v}, \mathbf{w}) = a \sum_{i \neq j} \frac{m_{i}m_{j}}{\|\mathbf{r}_{i} - \mathbf{r}_{j}\|^{a+2}}$$

$$\cdot \left(\frac{a+2}{\|\mathbf{r}_{i} - \mathbf{r}_{j}\|^{2}}(\mathbf{r}_{i} - \mathbf{r}_{j}, \mathbf{v}_{i} - \mathbf{v}_{j})(\mathbf{r}_{i} - \mathbf{r}_{j}, \mathbf{w}_{i} - \mathbf{w}_{j}) - (\mathbf{v}_{i} - \mathbf{v}_{j}, \mathbf{w}_{i} - \mathbf{w}_{j})\right)$$

$$+ b \sum_{i \neq j} \frac{m_{i}m_{j}}{\|\mathbf{r}_{i} - \mathbf{r}_{j}\|^{b+2}}$$

$$\cdot \left(\frac{b+2}{\|\mathbf{r}_{i} - \mathbf{r}_{j}\|^{2}}(\mathbf{r}_{i} - \mathbf{r}_{j}, \mathbf{v}_{i} - \mathbf{v}_{j})(\mathbf{r}_{i} - \mathbf{r}_{j}, \mathbf{w}_{i} - \mathbf{w}_{j}) - (\mathbf{v}_{i} - \mathbf{v}_{j}, \mathbf{w}_{i} - \mathbf{w}_{j})\right).$$

where $\mathbf{v}, \mathbf{w} \in \Omega$.

(3) The second derivative of the restriction $U: S^*_{I_0} \to \mathbb{R}$ is:

$$D^2 U/(S_{I_0}^*)(\mathbf{r})(\mathbf{v}, \mathbf{w}) = D^2 U(\mathbf{r})(\mathbf{v}, \mathbf{w}) + \frac{aW(\mathbf{r}) + bV(\mathbf{r})}{I_0} \langle \mathbf{v}, \mathbf{w} \rangle$$

Here (\cdot, \cdot) *denotes the usual inner product in* \mathbb{R}^2 *,* $\|\cdot\|$ *the norm in* \mathbb{R}^2 *, and I the moment of inertia. The same formulas are valid in* \mathbb{R} *,* \mathbb{R}^2 *and* \mathbb{R}^3 *.*

Proof. All the equations above can be derived by differentiating in local Cartesian coordinates. \Box

Now we can give a proof of Proposition 2. For $\mathbf{v}_i \in \mathbb{R}^2$, let $\mathbf{v}_i = (v_i^{\parallel}, v_i^{\perp})$ where $v_i^{\parallel} \in l$ and $v_i^{\perp} \in l^{\perp}$. Then we can write $\mathbf{v} = (\mathbf{v}^{\parallel}, \mathbf{v}^{\perp})$ with $\mathbf{v}^{\parallel} = (v_1^{\parallel}, \dots, v_n^{\parallel})$, $\mathbf{v}^{\perp} = (v_1^{\perp}, \dots, v_n^{\perp})$ for each $\mathbf{v} \in \Omega$. If $\mathbf{r} \in S_l \subset S_{l_0}$, $\mathbf{r} \notin \Delta$, we have $T_{\mathbf{r}}(S_{l_0}) = \{\mathbf{v} \in \Omega \mid \langle \mathbf{v}, \mathbf{r} \rangle = 0\}$ and $T_{\mathbf{r}}(S_l) = \{\mathbf{w} \in \Omega_l \mid \langle \mathbf{w}, \mathbf{r} \rangle = 0\}$ where, as usual, Ω is endowed with the mass scalar product. If $\mathbf{v} \in T_{\mathbf{r}}(S_{l_0})$ and $\mathbf{v} = (\mathbf{v}^{\parallel}, \mathbf{v}^{\perp})$, then $\mathbf{v}^{\parallel} \in \Omega_l$ and $\langle \mathbf{v}, \mathbf{r} \rangle = \langle \mathbf{v}^{\parallel}, \mathbf{r} \rangle$. Thus $\mathbf{v}^{\parallel} \in T_{\mathbf{r}}(S_l)$, because $\langle \mathbf{v}, \mathbf{r} \rangle = 0$ implies $\langle \mathbf{v}^{\parallel}, \mathbf{r} \rangle = 0$.

By Lemma 2 it follows that if $\mathbf{r} \in S_l \setminus \Delta$ and $\mathbf{v} \in T_{\mathbf{r}}(S_{l_0})$, then $DU(\mathbf{r})(\mathbf{v}) = DU(\mathbf{r})(\mathbf{v}^{\parallel})$. So $DU(\mathbf{r})(\mathbf{v}^{\parallel}) = 0$ implies that $DU(\mathbf{r})(\mathbf{v}) = 0$. This completes the proof of Proposition 2.

Lemma 3. $\mathbb{R}P^{n-2}$ has n!/2 components.

Proof. Let $\mathbf{r} = (\mathbf{r}_1, \dots, \mathbf{r}_n) \in S_l \setminus \Delta$ and let $\mathbf{r}_1 < \dots < \mathbf{r}_n \in \mathbb{R}$ (we use the fact that the \mathbf{r}_i are all distinct). Let $\alpha = (i_1, \dots, i_n)$ be an arbitrary permutation of the numbers $(1, 2, \dots, n)$. If we apply the permutation to the initial vector \mathbf{r} , we map it to a different component defined uniquely by the given permutation. Therefore the set $S_l \setminus \Delta$ has n! components and the quotient space $\mathbb{R}P^{n-2} \setminus \Delta$ has n!/2 components. \Box

We can now prove Moulton's theorem for quasihomogeneous potentials. By applying parts (2) and (3) of Lemma 2, we see that $D^2 U/(S_l \setminus \Delta)$ is a positive definite form, and consequently \tilde{U} is convex. This shows that \tilde{U} has a unique minimum in each component of $\mathbb{R}P^{n-2}$. Thus there are n!/2 critical points and hence n!/2 central configurations.

Remark 1. We have identified the symmetric central configurations, otherwise the number of classes of central configurations would be n!.

5. Planar central configurations

In this and subsequent sections, we will restrict our study to Manev-type quasihomogeneous potentials, namely those U for which a = 1 (see also [5]). They form an important class of quasihomogeneous potentials, derived from the Manev law, which can explain the perihelion advance of the planet Mercury within the framework of classical mechanics (for more details see [4] and [3]). Since for any planar central configuration in the Manev-type three-body problem, the mutual distances are geometrically independent, we can solve the equations defining the central configurations in terms of the mutual distances. To be precise, we state here a result whose proof can be found in [2].

Lemma 4. Let $u = f(\mathbf{x})$ be a function with $\mathbf{x} = (x_1, x_2, \dots, x_n)$, $x_1 = g_1(\mathbf{y})$, $x_2 =$ $g_2(\mathbf{y}), \ldots, x_n = g_n(\mathbf{y}), \mathbf{y} = (y_1, y_2, \ldots, y_m) \text{ and } m \ge n.$ If rank (A) = n, where

$$A = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \cdots & \frac{\partial x_n}{\partial y_1} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_1}{\partial y_m} & \cdots & \frac{\partial x_n}{\partial y_m} \end{pmatrix},$$
(13)

then $\nabla f(\mathbf{x}) = \mathbf{0}$ if and only if $\nabla u(\mathbf{y}) = \mathbf{0}$.

0

Let us now consider the three-body case, and for this purpose we will use the notation $\mathbf{r}_i = (q_{i1}, q_{i2})$ for i = 1, 2, 3. From Lemma 4 we have that if rank (A) = 3, where

$$A = \begin{pmatrix} \frac{\partial r_{12}}{\partial q_{11}} & \frac{\partial r_{13}}{\partial q_{11}} & \frac{\partial r_{23}}{\partial q_{11}} \\ \frac{\partial r_{12}}{\partial q_{12}} & \frac{\partial r_{13}}{\partial q_{12}} & \frac{\partial r_{23}}{\partial q_{12}} \\ \frac{\partial r_{12}}{\partial q_{21}} & \frac{\partial r_{13}}{\partial q_{21}} & \frac{\partial r_{23}}{\partial q_{21}} \\ \frac{\partial r_{12}}{\partial q_{22}} & \frac{\partial r_{13}}{\partial q_{21}} & \frac{\partial r_{23}}{\partial q_{21}} \\ \frac{\partial r_{12}}{\partial q_{22}} & \frac{\partial r_{13}}{\partial q_{22}} & \frac{\partial r_{23}}{\partial q_{22}} \\ \frac{\partial r_{12}}{\partial q_{31}} & \frac{\partial r_{13}}{\partial q_{31}} & \frac{\partial r_{23}}{\partial q_{32}} \\ \frac{\partial r_{12}}{\partial q_{32}} & \frac{\partial r_{13}}{\partial q_{31}} & \frac{\partial r_{23}}{\partial q_{32}} \end{pmatrix} = \begin{pmatrix} \frac{q_{11} - q_{21}}{r_{12}} & \frac{q_{11} - q_{31}}{r_{13}} & 0 \\ \frac{q_{12} - q_{22}}{r_{12}} & \frac{q_{12} - q_{32}}{r_{23}} \\ 0 & -\frac{q_{11} - q_{21}}{r_{12}} & 0 & \frac{q_{22} - q_{32}}{r_{23}} \\ 0 & -\frac{q_{11} - q_{31}}{r_{13}} & -\frac{q_{21} - q_{31}}{r_{23}} \\ 0 & -\frac{q_{12} - q_{32}}{r_{13}} & -\frac{q_{22} - q_{32}}{r_{23}} \end{pmatrix}$$

then $\nabla U(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3) = 0$ if and only if $\nabla U(r_{12}, r_{13}, r_{23}) = 0$.

Some straightforward computations show that the rank (A) = 3 if and only if

$$\det \begin{pmatrix} q_{11} & q_{12} & 1\\ q_{21} & q_{22} & 1\\ q_{31} & q_{32} & 1 \end{pmatrix} \neq 0.$$

This determinant is twice the oriented area of the triangle formed by the three particles. In short, if \mathbf{r}_1 , \mathbf{r}_2 and \mathbf{r}_3 are not collinear, then $\nabla U(r_{12}, r_{13}, r_{23}) = \mathbf{0}$ if and only if $\nabla U(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3) = 0$.

Using Lemma 4 in order to find the planar central configurations, we first need to solve the equation

$$\nabla U = \sigma \nabla I \tag{14}$$

in terms of the mutual distances r_{ij} , taking into account the fact that the moment of inertia, I, can be written in terms of the mutual distances as $I = (1/\tilde{m}) \sum_{i=1}^{n} m_i m_j r_{ij}^2$, where \tilde{m} is the total mass. So, for fixed *i* and *j*, we have

$$-\frac{m_im_j}{r_{ij}^2} - b\frac{m_im_j}{r_{ij}^{b+1}} = 2\frac{\sigma}{\tilde{m}}m_im_jr_{ij}$$

Multiplying by r_{ij}^{b+1} , we obtain

$$f(r_{ij}) := 2\sigma r_{ij}^{b+2} + \tilde{m} r_{ij}^{b-1} + \tilde{m} b = 0.$$

Regarding the above equation as a polynomial in the variable r_{ij} , since $\sigma < 0$, $f(0) = b\tilde{m} > 0$ and the coefficients polynomial have just one change of sign, we can verify easily that the function f has exactly one positive root. Observe that the function f only depends on the total mass \tilde{m} , and therefore the respective solution for $f(r_{ij})$ is the same for all mutual distances. We have thus proved the following result.

Theorem 2. In the Manev-type three-body problem, for any values of the masses, there are exactly two equilateral central configurations, which correspond to the two possible orientations of a triangle in a plane.

6. A framework for the study of collisions

We will further study the dynamics at an near total collision for Manev-type *n*-body problems. A convenient framework for this purpose is given by the so-called McGehee coordinates [8],

$$\rho = (\mathbf{r}^{t} M \mathbf{r})^{1/2}$$

$$\mathbf{s} = \rho^{-1} \mathbf{r}$$

$$v = \rho^{b/2} (\mathbf{p}^{t} \mathbf{s})$$

$$\mathbf{u} = \rho^{b/2} (\mathbf{p} - (\mathbf{p}^{t} \mathbf{s}) M \mathbf{s}),$$
(15)

where $M = \text{diag}(m_1, m_1, m_2, m_2, \dots, m_n, m_n)$, $\mathbf{r} = (\mathbf{r}_1, \dots, \mathbf{r}_n)$, and $\mathbf{p} = (\mathbf{p}_1, \dots, \mathbf{p}_n)$. After a reparametrization of the time variable,

$$\mathrm{d}\tau = r^{-1-b/2}\mathrm{d}t,\tag{16}$$

the equations of motion (5) become

$$\rho' = \rho v$$

$$v' = \frac{b}{2}v^{2} + \mathbf{u}^{t}M^{-1}\mathbf{u} - \rho^{b-1}W(\mathbf{s}) - bV(\mathbf{s})$$

$$\mathbf{s}' = M^{-1}\mathbf{u}$$

$$\mathbf{u}' = \left(\frac{b}{2} - 1\right)\mathbf{u}v - (\mathbf{u}^{t}M^{-1}\mathbf{u})M\mathbf{s} + \rho^{b-1}[W(\mathbf{s})M\mathbf{s} + \nabla W(\mathbf{s})] + bV(\mathbf{s})M\mathbf{s} + \nabla V(\mathbf{s}).$$
(17)

Here the prime denotes differentiation with respect to the new (fictitious) time variable τ , and the old notation is maintained for the new dependent variables, which are now functions of τ . Furthermore, the new variables fulfil the constraints $\mathbf{s}^t M \mathbf{s} = 1$ and $\mathbf{u}^t \mathbf{s} = 0$.

In these coordinates the energy integral (6) turns into the relation

$$\frac{1}{2}(\mathbf{u}^{t}M^{-1}\mathbf{u} + v^{2}) - \rho^{b-1}W(\mathbf{s}) - V(\mathbf{s}) = h\rho^{b}.$$
(18)

We define the total collision manifold as

$$C = \{(\rho, \mathbf{s}, v, \mathbf{u}) \mid \rho = 0, \mathbf{u}^{t} M^{-1} \mathbf{u} + v^{2} - 2V(\mathbf{s}) = 0\}.$$
(19)

Notice that $\rho' = 0$ if $\rho = 0$, so *C* (which is an analytic submanifold of codimension 1 in the boundary of the phase space) is invariant under the flow of the system (17). By continuity of the solutions with respect to initial conditions, the flow on *C* provides important information about the orbits close to triple collision (see [8] for more details). The total collision manifold can also be regarded as an invariant boundary pasted onto each energy surface:

$$E_{h} = \left\{ \left(\rho, v, \mathbf{s}, \mathbf{u} \right) \middle| \frac{1}{2} (\mathbf{u}^{t} M^{-1} \mathbf{u} + v^{2}) - \rho^{b-1} W(\mathbf{s}) - V(\mathbf{s}) = h \rho^{b} \right\}.$$
 (20)

These concepts are ideal for understanding the qualitative behaviour of total- and near-totalcollision solutions.

7. Collision and near-collision dynamics

In this section we will study the dynamics of total- and near-total-collision orbits of the Manev-type *n*-body problem. An important role in this study is played by central configurations and by the solutions that can be derived from them.

In the planar Newtonian *n*-body problem, a rigid rotation of a central configuration is called a relative equilibrium; in rotating coordinates, relative equilibria are fixed points. A non-rotating homothetic orbit of a central configuration is called a homothety. The composition of a relative equilibrium and a homothety is called a homographic solution.

In the Manev-type three-body problem, since the potential only depends of the bodies' mutual distances, the central configurations are invariant under rotations, so any central configuration determines a particular periodic orbit, which in a rotating frame is a fixed point. So in the Manev-type three-body problem, any central configuration corresponds to a relative equilibrium.

In the Newtonian case, any central configuration also corresponds to a homothetic orbit. But is this valid for Manev-type potentials too? As we will further prove (see Theorem 5 and Section 8), this property is not satisfied in general. To show this, and to determine under what circumstances homothetic solutions still exist, we will prove several preliminary results.

Notice that the flow on C is given by the equations

$$v' = \frac{b}{2}v^{2} + \mathbf{u}^{t}M^{-1}\mathbf{u} - bV(\mathbf{s})$$

$$\mathbf{s}' = M^{-1}\mathbf{u}$$

$$\mathbf{u}' = \left(\frac{b}{2} - 1\right)\mathbf{u}v - (\mathbf{u}^{t}M^{-1}\mathbf{u})M\mathbf{s} + bV(\mathbf{s})M\mathbf{s} + \nabla V(\mathbf{s}).$$
(21)

The equilibrium points of system (21) are given by $\mathbf{u} = 0$, $v = \pm \sqrt{2V(\mathbf{s})}$, where \mathbf{s} must be a critical point for the function $V(\mathbf{s})$ restricted to the unit sphere corresponding to the mass matrix M. The masses are involved because the equation $bV(\mathbf{s})M\mathbf{s} + \nabla V(\mathbf{s}) = 0$ must be satisfied. But these are the critical points of the function \tilde{V} , which is the restriction of the homogeneous potential V to the unit sphere given by the mass matrix M. Such critical points correspond to the central configurations of the homogeneous potential V.

Proposition 3. For any value of b > 2, the flow on the total collision manifold C is gradient-like with respect to the coordinate -v (i.e. the flow increases with respect to -v along nonequilibrium solutions).

Proof. The energy relation (18), restricted to *C*, takes the form

$$\mathbf{u}^t M^{-1} \mathbf{u} + v^2 - 2V(\mathbf{s}) = 0$$

Using the above expression and (21), we get that

$$v' = \left(1 - \frac{b}{2}\right) \mathbf{u}^t M^{-1} \mathbf{u} \tag{22}$$

on *C*. If $\mathbf{u} \neq \mathbf{0}$ then v' < 0 is increasing with respect to -v. On the other hand if $\mathbf{u} = \mathbf{0}$ then $\mathbf{u}' = bV(\mathbf{s})M\mathbf{s} + \nabla V(\mathbf{s})$, i.e. $\mathbf{u}' = 0$ only if \mathbf{s} is a critical point of \tilde{V} . Consequently -v is strictly increasing along nonequilibrium solutions, which means that the vector field is gradient-like with respect to -v. \Box

Denote by $ind(s_0)$ the index of the critical point s_0 , i.e. the number of eigenvalues of $D^2 \tilde{V}(s_0)$ with negative real part. Then we can prove the following result.

Theorem 3. Let \mathbf{s}_0 be a nondegenerate central configuration of the planar n-body problem with potential V and b > 2. Then the dimensions of $W^u(\mathbf{s}_0^+)$ and $W^s(\mathbf{s}_0^-)$ are the same and equal to $2n - 2 - \operatorname{ind}(\mathbf{s}_0)$ in E_h . The dimensions of $W^s(\mathbf{s}_0^+)$ and $W^u(\mathbf{s}_0^-)$ are the same and equal to $2n - 4 + \operatorname{ind}(\mathbf{s}_0)$ in E_h . The dimension of E_h is 4n - 5.

Proof. Let \mathbf{s}_0 be a central configuration, $v = \pm \sqrt{2V(\mathbf{s}_0)}$, and $\mathbf{u} = 0$, then the equation of motion restricted to E_h are

$$\rho' = \rho v$$

$$v' = \left(1 - \frac{b}{2}\right) \mathbf{u}^t M^{-1} \mathbf{u} + (b - 1)\rho^{b-1} W(\mathbf{s}) + bh\rho^b$$

$$\mathbf{s}' = M^{-1} \mathbf{u}$$

$$\mathbf{u}' = \left(\frac{b}{2} - 1\right) \mathbf{u} v - (\mathbf{u}^t M^{-1} \mathbf{u}) M \mathbf{s} + \rho^{b-1} [W(\mathbf{s}) M \mathbf{s} + \nabla W(\mathbf{s})] + bV(\mathbf{s}) M \mathbf{s} + \nabla V(\mathbf{s}).$$
(23)

Taking into account the centre of mass and linear momentum integrals as well as the restrictions $\mathbf{s}^t M \mathbf{s} = 1$ and $\mathbf{u}^t \mathbf{s} = 0$ of the McGehee coordinates, the above system has dimension 4n - 4.

Linearizing the system, the eigenvalues for b > 2 are given by the matrix equation

$$\begin{pmatrix} v & 0 & 0 & \dots & \dots & 0 \\ 0 & 0 & * & \dots & & * \\ \vdots & \vdots & O_{2n-3} & I_{2n-3} \\ \vdots & \vdots & A & \left(\frac{b}{2} - 1\right) v I_{2n-3} \\ 0 & 0 & & & \end{pmatrix} - \mu I_{4n-4} = O_{4n-4},$$
(24)

where I_N is the $N \times N$ identity matrix, O_N is the $N \times N$ zero matrix, A denotes the Hessian matrix of \tilde{V} (i.e. the potential restricted to the sphere of constant moment of inertia) and * denotes an element without importance in the computation of the eigenvalues.

It is clear that the first two eigenvalues are $v \neq 0$ (since $V(\mathbf{s}_0) \neq 0$) and 0. To obtain the remaining eigenvalues of Eq. (24), suppose \mathbf{z} is a (2n - 3)-vector satisfying

$$A\mathbf{z} = \lambda_i \mathbf{z} \tag{25}$$

for i = 1, ..., 2n - 3, *i* fixed, where $\lambda_1, ..., \lambda_{2n-3}$ are the eigenvalues of *A*. Then

$$\begin{pmatrix} O_{2n-3} & I_{2n-3} \\ A & (b/2-1)vI_{2n-3} \end{pmatrix} \begin{pmatrix} \mathbf{z} \\ \mu \mathbf{z} \end{pmatrix} = \begin{pmatrix} \mu \mathbf{z} \\ \{\lambda_i + (b/2-1)v\mu\}\mathbf{z} \end{pmatrix}$$

Consequently μ is a root of Eq. (24) if

$$\mu^2 - (b/2 - 1)v\mu - \lambda_i = 0,$$

which gives

$$\mu_i^{1,2} = \frac{1}{4} \left\{ (b-2)v \pm \sqrt{(2-b)^2 v^2 + 16\lambda_i} \right\}$$

for i = 1, ..., 2n - 3.

Then if $v = \sqrt{V(\mathbf{s}_0)}$, the differential matrix of the vector field restricted to E_h has $2n - 2 - \operatorname{ind}(\mathbf{s}_0)$ eigenvalues with positive real part and $2n - 4 + \operatorname{ind}(\mathbf{s}_0)$ with negative real part. The values of the dimensions are switched if $v = -\sqrt{V(\mathbf{s}_0)}$. \Box

Theorem 4. Let \mathbf{s}_0 be a central configuration of the collinear *n*-body problem with potential *V* and b > 2. Then the dimensions of $W^u(\mathbf{s}_0^+)$ and $W^s(\mathbf{s}_0^-)$ are the same and equal to n - 1 in E_h . The dimensions of $W^s(\mathbf{s}_0^+)$ and $W^u(\mathbf{s}_0^-)$ are the same and equal to n - 2 in E_h . The dimension of E_h is 2n - 3.

Proof. The proof is similar to the one of the previous theorem. Let \mathbf{s}_0 be a central configuration, $v = \pm \sqrt{2V(\mathbf{s}_0)}$. The equations of motion restricted to E_h are given by Eq. (23), with the obvious modifications.

Linearizing the system, the eigenvalues in the case b > 2 are given by the following matrix equation

$$\begin{pmatrix} v & 0 & 0 & \dots & \dots & 0 \\ 0 & 0 & * & \dots & & * \\ \vdots & \vdots & O_{n-2} & I_{n-2} \\ \vdots & \vdots & A & \left(\frac{b}{2} - 1\right) v I_{n-2} \\ 0 & 0 & & & \end{pmatrix} - \mu I_{2n-2} = O_{2n-2},$$
(26)

where I_N and O_N are defined as before. Again A is the Hessian matrix of \tilde{V} and * denotes an element without importance in the computation of the eigenvalues. The first two eigenvalues are $v \neq 0$ (since $V(\mathbf{s}_0) \neq 0$) and 0. If $\lambda_1, \ldots, \lambda_{n-2}$ be the eigenvalues of A, then

$$\mu_i^{1,2} = \frac{1}{4} \left\{ (b-2)v \pm \sqrt{(2-b)^2 v^2 + 16\lambda_i} \right\}$$

for i = 1, ..., n-2. Note that, in this case, Lemma 4 implies that A is positive definite, and thus the eigenvalues $\lambda_1, ..., \lambda_{n-2}$ are all positive. Consequently, for v > 0, μ_i^1 is negative and μ_i^2 positive. However, for v < 0, μ_i^1 is positive, whereas μ_i^2 is negative. This concludes the proof.

We will further state and prove a result that clarifies under what circumstances homothetic solutions exist.

Theorem 5. A solution of the Manev-type n-body problem is homothetic if and only if the particles form, at all times, a simultaneous central configuration for the potentials V and W.

Proof. Assume that the solution is homothetic, then $\mathbf{s} \equiv \mathbf{s}_0$, where \mathbf{s}_0 is a constant. Therefore $\mathbf{s}' \equiv 0$ and, from the second of Eq. (21), $\mathbf{u} \equiv 0$. Thus the homothetic orbits are confined to the invariant plane

$$\mathcal{P} = \{(\rho, \mathbf{s}, v, \mathbf{u}) \mid \mathbf{s} = \mathbf{s}_0, \mathbf{u} = \mathbf{0}\}.$$
(27)

So $\mathbf{u}' = 0$ implies that $\rho^{b-1}[W(\mathbf{s}_0)M\mathbf{s}_0 + \nabla W(\mathbf{s}_0)] + bV(\mathbf{s}_0)M\mathbf{s}_0 + \nabla V(\mathbf{s}_0) = 0$. If ρ is not constant then there are $\rho_1 \neq 0$ and $\rho_2 \neq 0$ with $\rho_1 \neq \rho_2$ such that

$$\rho_1^{b-1}[W(\mathbf{s}_0)M\mathbf{s}_0 + \nabla W(\mathbf{s}_0)] = -[bV(\mathbf{s}_0)M\mathbf{s}_0 + \nabla V(\mathbf{s}_0)]$$

$$\rho_2^{b-1}[W(\mathbf{s}_0)M\mathbf{s}_0 + \nabla W(\mathbf{s}_0)] = -[bV(\mathbf{s}_0)M\mathbf{s}_0 + \nabla V(\mathbf{s}_0)].$$
(28)

This means that $[bV(\mathbf{s}_0)M\mathbf{s}_0 + \nabla V(\mathbf{s}_0)] = 0$ and $[W(\mathbf{s}_0)M\mathbf{s}_0 + \nabla W(\mathbf{s}_0)] = 0$, i.e. that \mathbf{s}_0 is a simultaneous central configuration for the potentials V and W. If ρ is constant, $\rho' = 0$ and either $\rho \equiv 0$ or $v \equiv 0$. The first case is trivial, whereas in the latter case $-\rho^{b-1}W(\mathbf{s}) - bV(\mathbf{s}) = 0$. But this is impossible since V > 0 and W > 0.

If $\mathbf{s} \equiv \mathbf{s}_0$ is, at all times, a simultaneous central configuration for *V* and *W*, then the solution is obviously homothetic. \Box

The next result proves the existence and uniqueness of heteroclinic homothetic solutions for b > 1.

Theorem 6. Let \mathbf{s}_0 be a simultaneous central configuration for the potentials V and W. Then, if b > 1, every energy surface of negative constant (h < 0), contains a unique homothetic solution defined on $(-\infty, \infty)$, satisfying $\mathbf{s} = \mathbf{s}_0$ for all times and such that $\rho(\tau) \to 0$ when $\tau \to \pm \infty$. In other words the solution begins and ends in a total collapse, maintaining for all times the same central configuration.

1436

Proof. Since \mathbf{s}_0 is a simultaneous central configuration for the potentials *V* and *W*, we have that $[bV(\mathbf{s}_0)M\mathbf{s}_0 + \nabla V(\mathbf{s}_0)] = 0$ and $[W(\mathbf{s}_0)M\mathbf{s}_0 + \nabla W(\mathbf{s}_0)] = 0$. Consequently, the set

$$\mathcal{P} = \{(\rho, \mathbf{s}, v, \mathbf{u}) \mid \mathbf{s} = \mathbf{s}_0, \mathbf{u} = \mathbf{0}\}$$

is invariant for the equations of motion. Restricting these equations to \mathcal{P} , we get

$$\rho' = \rho v v' = \frac{b}{2}v^2 - \rho^{b-1}W(\mathbf{s}_0) - bV(\mathbf{s}_0),$$
(29)

while the energy relation becomes

$$\frac{1}{2}v^2 - \rho^{b-1}W(\mathbf{s}_0) - V(\mathbf{s}_0) = h\rho^b.$$

Eq. (29) become

$$\rho' = \rho v$$

$$v' = (b-1)\rho^{b-1}W(\mathbf{s}_0) + b\rho^b h.$$

This leads to

$$\frac{\mathrm{d}v}{\mathrm{d}\rho} = \frac{1}{v} [(b-1)\rho^{b-2}W(\mathbf{s}_0) + b\rho^{b-1}h],$$

which yields

$$\frac{v^2}{2} = \rho^{b-1} W(\mathbf{s}_0) + \rho^b h + K,$$
(30)

where, if b > 1, we choose $K = V(\mathbf{s}_0)$. If b > 1 and $h \ge 0$, then $|v| \ge \pm \sqrt{2V(\mathbf{s}_0)}$ and the homothetic orbits are not heteroclinic. If h < 0, there is a unique curve connecting the points $(\sqrt{2V(\mathbf{s}_0)}, 0)$ and $(-\sqrt{2V(\mathbf{s}_0)}, 0)$ on the plane (v, ρ) . These facts prove the theorem. \Box

The following result shows that the above property is also true for the equilateral central configurations.

Corollary 1. Let \mathbf{s}_0 be an equilateral central configuration for the potential U. Then, if b > 1, every energy surface of negative constant (h < 0) contains a unique heteroclinic homothetic solution.

Proof. Clearly \mathbf{s}_0 is a simultaneous central configuration for the potentials *V* and *W*. The proof follows from Theorem 6. \Box

Two submanifolds \mathcal{E}_1 and \mathcal{E}_2 of a submanifold \mathcal{E} are said to be transverse at a point x if one of the following situation arises:

(1) £₁ ∩ £₂ = Ø;
(2) x ∈ £₁ ∩ £₂ and T_xE₁ + T_xE₂ = T_xE, where T_xE denotes the tangent space to E at the point x.

We can now prove the following result:

Theorem 7. In the planar Manev-type n-body problem with b > 2, a necessary condition for having a transversal homothetic solution $\gamma_h(\mathbf{s}_0)$ in E_h with h < 0 is that \tilde{V} be a nondegenerate minimum at the point \mathbf{s}_0 associated with the homothetic solution.

Proof. Let $\gamma_h(\mathbf{s}_0)$ be a transversal homothetic solution in E_h with h < 0. Then by Theorem 3 both $W^u(\mathbf{s}_0^+)$ and $W^s(\mathbf{s}_0^-)$ are $(2n - 2 - \operatorname{ind}(\mathbf{s}_0))$ -dimensional and E_h is (4n - 5)-dimensional. Since $\gamma_h(\mathbf{s}_0) \in W^u(\mathbf{s}_0) \cap W^s(\mathbf{s}_0)$ and $\gamma_h(\mathbf{s}_0)$ is transversal we have that

$$\dim E_h \leq \dim W^u(\mathbf{s}_0) + \dim W^s(\mathbf{s}_0) - 1$$

That is $4n-5 \le 4n-5-2$ ind(\mathbf{s}_0). Therefore ind(\mathbf{s}_0) = 0 and the function \tilde{V} has a nondegenerate minimum at \mathbf{s}_0 . \Box

8. Simultaneous configurations and relative equilibria

In this closing section, we will show that, for most choices of the masses in the quasihomogeneous three-body problem, the collinear central configurations of the potential U are not simultaneous relative equilibria for V and W.

Theorem 8. Let Σ_3 be the set of masses $(m_1, m_2, m_3) \in \mathbb{R}^3_+$ for which the collinear configurations are simultaneous central configurations for the potentials V and W. Then the set Σ_3 is nonempty and nowhere dense in \mathbb{R}^3_+ .

Proof. Assume the configuration $\mathbf{s}_V(m_1, m_2, m_3)$ is a collinear central configuration for V and $\mathbf{s}_W(m_1, m_2, m_3)$ is a collinear central configuration for W. In [7], Euler found a complicated formula that expresses the ratio of the distances between the masses for any rectilinear central configuration in the Newtonian case. Euler's formula can be directly extended to any homogeneous potential. Moreover, the fact that Euler's expression is an analytic function of the masses remains true in the homogeneous case. Therefore both \mathbf{s}_V and \mathbf{s}_W are analytic functions of m_1, m_2 and m_3 , as long as the masses are positive. Consequently the function $z = \mathbf{s}_V - \mathbf{s}_W$ is also an analytic function of the masses.

For the function V, Euler's formula depends on a, whereas for W it depends on b. So in general $\mathbf{s}_V \neq \mathbf{s}_W$, therefore for every a and b with $a \neq b$ there are values of the masses for which $z \neq 0$. Since z is a nonzero analytic function, its zeroes form a nowhere dense set.

The nonemptiness of the set of simultaneous central configurations follows from noticing that if $m_1 = m_3$ and the mass m_2 is located halfway between the other two, then the three masses form a simultaneous central configuration for V and W.

A consequence of Theorems 5 and 8 is that, for most values of the masses, there are no rectilinear homothetic orbits. More precisely:

Corollary 2. If $(m_1, m_2, m_3) \in \mathbb{R}^3_+ \setminus \Sigma_3$, then there are no rectilinear homothetic orbits.

This shows that the rectilinear homothetic orbits are characteristic to homogeneous potentials, but they prove unlikely in the quasihomogeneous case.

References

- [1] R. Abraham, J.E. Marsden, Foundation of Mechanics, 2nd ed., Benjamin, New York, 1978.
- [2] M. Corbera, J. Llibre, E. Pérez-Chavela, Equilibrium points and central configurations for the Lennard-Jones 2- and 3-body problems, Celestial Mechanics and Dynamical Astronomy 89 (3) (2004) 235–266.
- [3] S. Craig, F. Diacu, E.A. Lacomba, E. Pérez-Chavela, On the anisotropic Manev problem, Journal of Mathematical Physics 40 (1999) 1–17.
- [4] F. Diacu, Near-collision dynamics for particle systems with quasihomogeneous potentials, Journal of Differential Equations 128 (1996) 58–77.

- [5] F. Diacu, V. Mioc, C. Stoica, Phase-space structure and regularization of Manev-type problems, Nonlinear Analysis 41 (2000) 1029–1055.
- [6] F. Diacu, E. Pérez-Chavela, M. Santoprete, Saari's conjecture in the collinear case, Transactions of the American Mathematical society 357 (10) (2005) 4215–4223.
- [7] L. Euler, De motu rectilineo trium corporum se mutuo attrahentium, Novi Commentarii Academiae Scientarum Petropolitanae 11 (1767) 144–151.
- [8] R. McGehee, Triple collision in the collinear three-body problem, Inventiones Mathematicae 27 (1974) 191–227.
- [9] K. Meyer, G. Hall, Introduction to Hamiltonian Dynamical Systems and the n-Body Problem, in: Applied Mathematical Science, vol. 90, Springer-Verlag, 1992.
- [10] F.R. Moulton, The straight line solutions of the problem of N bodies, Annals of Mathematics 12 (1910) 1–17.
- [11] E. Pérez-Chavela, D. Saari, A. Susin, Z. Yan, Central configurations in the charged three body problem, Contemporary Mathematics 198 (1996) 137–155.
- [12] E. Pérez-Chavela, L. Vela-Arévalo, Triple collision in the quasihomogeneous collinear three-body problem, Journal of Differential Equations 148 (1998) 186–211.
- [13] D. Saari, On the role and properties of n-body central configurations, Celestial Mechanics 21 (1980) 9–20.
- [14] S. Smale, Problems on the nature of relative equilibria in celestial mechanics, Lecture Notes in Mathematics 197 (1971) 194–198.
- [15] S. Smale, Mathematical problems for the next century, Mathematical Intelligencer 20 (2) (1998) 7–15.
- [16] A. Wintner, The Analytical Foundations of Celestial Mechanics, Princeton University Press, Princeton, NJ, 1941.